

A stratification of the space of all k -planes in \mathbb{C}^n

Allen Knutson (Cornell)

AMS-MAA talk, Boston joint meetings, 2012

Abstract

To each $k \times n$ matrix M of rank k , we associate a *juggling pattern* of periodicity n with k balls. The juggling pattern actually only depends on the k -plane spanned by the rows, so gives a decomposition of the “Grassmannian” of all k -planes in n -space.

There are many connections between the geometry and the juggling. For example, the natural topology on the space of matrices induces a partial order on the space of juggling patterns, which indicates whether one pattern is “more excited” than another.

This same decomposition turns out to naturally arise from totally positive geometry [Lusztig 1994, Postnikov ~2004], characteristic p geometry [Knutson-Lam-Speyer 2011], and noncommutative geometry [Brown-Goodearl-Yakimov 2005]. It also arises by projection from the manifold of full flags in n -space, where there is no cyclic symmetry.

A discrete invariant of matrices.

For the purposes of this talk, an **invariant of matrices** is a function $f : \{\text{matrices}\} \rightarrow \text{somewhere}$ that is invariant under row operations, or equivalently, $f(M) = f(AM)$ for A invertible. One of the best known is $\text{rank} : M_{k \times n} \rightarrow \mathbb{N}$ (which is also invariant under column operations).

Today's is the following. Think of M as a list $\vec{v}_1, \dots, \vec{v}_n$ of k -dimensional column vectors, and extend it to be an infinite but periodic list, $\vec{v}_i = \vec{v}_{n+i}$. Then define

$$J_M : \mathbb{Z} \rightarrow \mathbb{Z}, \quad J_M(i) := \min \{j \geq i : \vec{v}_i \in \text{span}(\vec{v}_{i+1}, \dots, \vec{v}_j)\} \leq i + n.$$

For example,

$$\begin{array}{cccccccccc}
 \mathbf{v} & \mathbf{v} & \mathbf{v} & \mathbf{v} & \mathbf{v} & \mathbf{v} & \mathbf{v} & \mathbf{v} & \mathbf{v} & \mathbf{v} \\
 -1 & 0 & 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 \\
 \dots & 0 & 0 & \boxed{1} & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\
 & 0 & 0 & 0 & \boxed{1} & 1 & 0 & 0 & 0 & 0 & 1 & \dots \\
 & 1 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 0 & 0
 \end{array}$$

$J(i)-i:$
→ 6
→ 1
→ 5
→ 1
→ 5
→ 0

A nonobvious property: J_M is 1:1 and onto! What else is true about these J_M ?

Bounded juggling patterns, with a fixed periodicity n .

An **affine permutation** $J : \mathbb{Z} \rightarrow \mathbb{Z}$ is a function that's 1:1 and onto, with the periodicity $J(i + n) = J(i) + n \quad \forall i$. These form a group isomorphic to $S_n \ltimes \mathbb{Z}^n$, where $S_n := \text{Sym}(\mathbb{Z}/n)$ is the finite permutation group.

If we try to interpret $i \mapsto J(i)$ as “A ball thrown at time i comes down at time $J(i) - \frac{1}{2}$, and is then thrown at time $J(i)$ ” we had better insist $J(i) \geq i$, so balls land *after they are thrown*. Call such affine permutations **juggling patterns**. The number of balls in the air at time $i + \frac{1}{2}$, $\#\{k < i + \frac{1}{2} : J(k) > i + \frac{1}{2}\}$, is finite and (thankfully) independent of i .

What jugglers actually make use of is not J , but its associated **siteswap** $J(1)-1 \ J(2)-2 \ \dots \ J(n)-n$, the list of throw heights durations durations+ $\frac{1}{2}$. Useful theorem to come: the number of balls is the average of the siteswap.

Some examples: 3 ~ 3333, 4, 1, 51, 441, 4413, 330, 4440, 42, 552, 51414, 53...

If you want to see another hour of this, look up “knutson juggling” on YouTube.

Define a **bounded juggling pattern** to be an affine permutation J that not only satisfies $J(i) \geq i$, but also $J(i) \leq i + n$, for all i .

Theorem [Postnikov ~2004, juggling interpretation in K-Lam-Speyer 2011]. Each J_M (from the last page) is a bounded juggling pattern, and every k -ball period- n bounded juggling pattern arises from some $k \times n$ matrices of rank k .

Total positivity of matrices.

Matrices with real entries in which every submatrix has nonnegative determinant have been studied since the 1930s and impact many areas (see the entire book [Karlin 1968]). In our context we consider real $k \times n$ matrices where every $k \times k$ submatrix has determinant ≥ 0 . These have a surprising cyclic property, that will connect to the periodicity of our patterns:

Lemma. If $[\vec{v}_1 \cdots \vec{v}_n]$ is a totally nonnegative matrix, so is $[\vec{v}_2 \cdots \vec{v}_n \ (-1)^{k-1} \vec{v}_1]$.

These $\binom{n}{k}$ many $k \times k$ determinants are not independent; e.g. in 2×4 they satisfy

$$p_{13} p_{24} = p_{12} p_{34} + p_{14} p_{23}, \quad p_{ij} := \det(\text{columns } i \text{ and } j)$$

which is very stringent if we also require each $p_{ij} \geq 0$!

Theorem [Postnikov ~2004]. Let $B(M) = \{S \subseteq \{1, \dots, n\} : |S| = k, p_S \neq 0\}$, the **bases of the matroid** associated to the matrix M .

If M is totally nonnegative and rank k , then $B(M)$ and J_M determine each other, and $B(M)$ is called a **positroid**. (If $\text{rank}(M) \neq k$, then $B(M) = \emptyset$.)

The **positroid $\mathbb{R}_{\geq 0}$ -stratum** of totally nonnegative matrices with a given J_M is (nonempty and) homeomorphic to an open ball.

If one drops the total-nonnegativity assumption, the topology of a matroid stratum can be, in some senses, arbitrarily bad (Mnëv's universality theorem).

The Freshman's Dream, and splitting the Frobenius morphism.

Let R be a commutative ring in which $1 + 1 + \dots + 1 = 0$, added up p times. If R has no zero divisors, then p must be prime. We assume p is prime and say that R has **characteristic p** .

The Freshman's Dream. In a ring of characteristic p , $(a + b)^p = a^p + b^p$, i.e. $r \mapsto r^p$ is an endomorphism called the **Frobenius**.

Call an abelian group homomorphism $\varphi : R \rightarrow R$ a **Frobenius splitting** if

- $\varphi(r^p) = r, \forall r \in R$ so, φ is a one-sided inverse
- $\varphi(r^p q) = r \varphi(q)$ another desirable property of such a “ p th root” map.

Example. Let $R = \mathbb{F}_p[x]$, $\varphi(cx^k) = cx^{k/p}$ if $p \mid k$, 0 otherwise.

A similar rule works for $R = \mathbb{F}_p[x_1, \dots, x_n]$, or that modulo any monomial ideal, and many other φ exist for these R .

Example. Let $R = \mathbb{F}_p[a^2, a^3] \leq \mathbb{F}_p[a]$, so $R \cong \mathbb{F}_p[x, y]/\langle y^2 - x^3 \rangle$. Then $\nexists \varphi$.

It's easy to show that if R has a Frobenius splitting φ , then R must have no nilpotents. As the second example shows, though, the condition is much more stringent.

Compatibly split ideals.

In the category of “Frobenius split rings (R, φ) of characteristic p ” the right notion of ideal $I \leq R$ is one such that $\varphi(I) \leq I$, called a **compatibly split ideal**.

Theorem [Enescu–Hochster 2008, Schwede 2009, Kumar–Mehta 2009].

If R is a Frobenius split Noetherian ring (or more generally a Noetherian scheme with a Frobenius splitting on its structure sheaf), then it has only finitely many compatibly split ideals (resp. ideal sheaves).

Sad proposition [K]. If $R = \mathbb{F}_p[x_{11}, \dots, x_{kn}]$ is the functions on the space of $k \times n$ matrices, and $A = p_{12\dots k} p_{23\dots k+1} p_{34\dots k+2} \cdots p_{n-1\ n\ 12\dots k-2} p_{n12\dots k-1}$, then for $n, k > 1, n \neq k$ there is no splitting φ that compatibly splits $\langle A \rangle$.

Luckily we don’t want to apply this technology to *matrices*, but to rank k matrices up to row-equivalence. So some k columns $S \subseteq \{1, \dots, n\}$ must form a basis, and we can use up the row operations making them the identity matrix.

Theorem [K-Lam-Speyer 2011]. Let R_S be the functions on the (affine) space of $k \times n$ matrices whose columns S are an identity matrix. Then there is a unique splitting on R_S that compatibly splits the $\langle A \rangle$ above, and its compatibly split prime ideals are exactly given by the positroid stratification.

This is more cleanly stated as being about a splitting on the **Grassmannian of k -planes in n -space**, which has an atlas given by these $\binom{n}{k}$ affine patches.

A noncommutative deformation of the Grassmannian.

Let R be a vector space, and $\cdot_\epsilon : R \times R \rightarrow R$ a family of associative products on it, one for each number ϵ . If \cdot_0 is commutative, then we can think of (R, \cdot_0) as the ring of functions on a space $\text{Spec}(R, \cdot_0)$.

If $I \leq R$ is an ideal for every \cdot_ϵ , then it is for \cdot_0 , and defines a subset of $\text{Spec}(R, \cdot_0)$. But very few ideals arise this way, as noncommutative rings have far fewer of them than commutative rings do! One says that very few subvarieties “survive deformation to a noncommutative space”.

$R = \mathbb{C}[x_{11}, \dots, x_{kn}]$ has a family of products \cdot_ϵ described to first order by

$$x_{ij} \cdot_\epsilon x_{kl} = x_{kl} \cdot_\epsilon x_{ij} + \epsilon \text{sign}(k - i) \text{sign}(l - j) x_{il} x_{kj} + O(\epsilon^2)$$

Theorem [Brown-Goodearl-Yakimov 2006]. Let $I \leq R$ be a prime ideal of every (R, \cdot_ϵ) , invariant under scaling the columns ($x_{ij} \mapsto t_j x_{ij}$). Then $I \leq (R, \cdot_0)$ defines one of our positroid strata, and each stratum arises this way from a unique I .

(This is connected to the Frobenius splitting, as follows. The first-order term above defines a *Poisson 2-tensor*, which wedged with some column-scaling vector fields gives an *anticanonical tensor*. From that tensor one can build a map $\phi : R \rightarrow R$, which may or may not be a splitting; in this case it is.)

An application of the positroid stratification to juggling.

Let $J, J' : \mathbb{Z} \rightarrow \mathbb{Z}$ be two juggling patterns. Call J' a **simple excitation** of J if

- $J(i) = J'(i)$ unless $i \equiv a, b \pmod n$ for some pair $a < b$
- $J(a) < J(b)$ and $J'(a) = J(b), J'(b) = J(a)$
- for all c in the open interval (a, b) , $J(c) \notin (J(a), J(b))$.

Call J' an **excitation** of J if they are connected by a sequence of simple such. It is easy to see that J, J' must have the same number of balls, and their siteswaps must have the same average. Example (with a, b underlined):

$$\underline{5}1414 \succ 24\underline{4}14 \succ 24\underline{2}34 \succ 23334 \sim 333\underline{4}2 \succ 33333$$

Proposition. The unique least excited pattern with k balls is $J(i) = i + k$, with all throws being k s. There are $\binom{n}{k}$ most excited bounded juggling patterns with k balls, with $(n - k)$ 0-throws and k n -throws.

Corollary (stated before): the average of the siteswap is the number of balls.

Theorem [K-Lam-Speyer 2011]. The positroid stratum for J' is in the closure of the stratum for J if and only if J' is an excitation of J .

Jugglers had already known about the $b = a + 1$ simple excitations, but not these more general ones, nor that there is a well-defined **excitation number** given by the codimension of the corresponding stratum.

References

[Karlin 1968] S. Karlin, Total positivity, Stanford University Press, 1968.

[Lusztig 1994] G. Lusztig, Total positivity in reductive groups, Lie Theory and Geometry, Progress in Math., vol. 123, Birkhäuser Boston, 1994, pp. 531–568.

[Postnikov ~2004] A. Postnikov, Total positivity, Grassmannians, and networks, preprint. <http://arxiv.org/abs/math.CO/0609764>

[Brown-Goodearl-Yakimov 2005] K. A. Brown, K. A. Goodearl, and M. Yakimov, Poisson structures on affine spaces and flag varieties. I. Matrix affine Poisson space, Advances in Math., <http://arxiv.org/abs/math.QA/0501109>

[Enescu–Hochster 2008] F. Enescu and M. Hochster, The Frobenius structure of local cohomology, Algebra and Number Theory 2 (2008), no. 7, 721–754. <http://arxiv.org/abs/0708.0553>

[Schwede 2009] K. Schwede, F-adjunction, Algebra and Number Theory. Vol 3, no. 8, 907–950. (2009) <http://arxiv.org/abs/0901.1154>

[Kumar–Mehta 2009] V. B. Mehta and S. Kumar: Finiteness of the number of compatibly-split subvarieties, IMRN vol. 2009 (no. 19), 3595–3597. <http://arxiv.org/abs/math.QA/0901.2098>

[K-Lam-Speyer 2011] A.K., T. Lam, D. E Speyer, Projected Richardson varieties, to appear in Crelle’s journal. <http://arxiv.org/abs/1008.3939>